


Applicants: Timothy Norris et al.
Serial No.: 10/711,272
Filed: November 9, 2000
Page 6

No fee is deemed necessary in connection with the filing of this Supplemental Amendment and Supplemental Information Disclosure Statement. However, if any fee is required, authorization is hereby given to charge the amount of any such fee to Deposit Account No. 03-3125.

Respectfully submitted,

A handwritten signature in dark ink, appearing to read "Gary J. Gershik", is written over a horizontal line.

John P. White
Registration No. 28,678
Gary J. Gershik
Registration No. 39,992
Attorneys for Applicants
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number
WO 01/34574 A1

(51) International Patent Classification: C07D 239/04,
A61K 31/505

RAGGON, Jeffrey, W.; 48 Teecomwas Drive, Uncasville,
CT 06382 (US). SILBERMAN, Sandra, L.; 10 Heather
Lane, Randolph, NJ 07869 (US).

(21) International Application Number: PCT/US00/31009

(74) Agent: WHITE, John, P.; Cooper & Dunham LLP, 1185
Avenue of the Americas, New York, NY 10036 (US).

(22) International Filing Date:
9 November 2000 (09.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/164,907 11 November 1999 (11.11.1999) US
60/193,191 30 March 2000 (30.03.2000) US
60/206,420 23 May 2000 (23.05.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: OSI PHARMACEUTICALS, INC.
[US/US]; 106 Charles Lindbergh Boulevard, Union-
dale, NY 11553-3649 (US).

(72) Inventors: CONNELL, Richard, D.; 23 Jeremy Drive,
East Lyme, CT 06333 (US). MOYER, James, D.; 5 Jeff-
erson Drive, East Lyme, CT 06333 (US). MORIN, Michael,
J.; 8 Goundry Drive, Waterford, CT 06385 (US). KA-
JLI, Shama, M.; 190 Mistuxet Avenue, Mystic, CT 06355
(US). FOSTER, Barbara, A.; 31 Rossie Street, Mystic,
CT 06355 (US). FERRANTE, Karen, J.; 10 Adirondack
Drive, East Greenwich, RI 02818 (US). NORRIS, Tim-
othy; 27 Friar Tuck Drive, Gales Ferry, CT 06335 (US).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: STABLE POLYMORPH OF N-(3-ETHYNYLPHENYLAMINO)-6,7-BIS(2-METHOXYETHOXY)-4-QUINAZOLI-
NAMINE HYDROCHLORIDE, METHODS OF PRODUCTION, AND PHARMACEUTICAL USES THEREOF

(57) Abstract: The present invention relates to a stable crystalline form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quina-
zolinamine hydrochloride designated the B polymorph, its production in essentially pure form, and its use. The invention also relates
to the pharmaceutical compositions containing the stable polymorph B form of N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-
quinazolinamine as hydrochloride, as well as other forms of the compound, and to methods of treating hyperproliferative disorders,
such as cancer, by administering the compound.

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 8

WO 01/34574 A1

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 837 063 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
22.04.1998 Bulletin 1998/17

(21) Application number: 97307724.1

(22) Date of filing: 01.10.1997

(51) Int. Cl.⁵: C07D 403/12, A61K 31/505,
C07D 239/88, C07D 401/14,
C07D 401/06
// (C07D403/12, 239:00,
209:00)

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV RO SI

(30) Priority: 17.10.1996 US 28881 P


(71) Applicant: PFIZER INC.
New York, N.Y. 10017 (US)

(72) Inventors:
• Arnold, Lee Daniel
Westborough, Massachusetts 01581 (US)
• Sobolov-Jaynes, Susan Beth
Ivoryton, Connecticut 06442 (US)

(74) Representative:
Hayles, James Richard et al
Pfizer Limited,
Patents Department,
Ramsgate Road
Sandwich Kent CT13 9NJ (GB)

(54) 4-Aminoquinazoline derivatives

(57) This invention relates to certain 4-aminoquinazoline derivatives of the formula



and their pharmaceutically acceptable salts wherein R¹, Q¹, m, n, and Z are defined as in the specification. The compounds of formula I and pharmaceutically acceptable salts are useful for the treatment of hyperproliferative disorders and conditions in mammals.

EP 0 837 063 A1

PRODUCTION OF AMINOPHENYLACETYLENE COMPOUND

Patent Number: JP10036325
Publication date: 1998-02-10
Inventor(s): YAMAKAWA KAZUYOSHI; SATO TADAHISA
Applicant(s): FUJI PHOTO FILM CO LTD
Requested Patent: ☐ JP10036325
Application Number: JP19960207786 19960718
Priority Number(s):
IPC Classification: C07C211/45; C07C209/36; C07C213/02; C07C215/68; C07C215/70; C07F7/10
EC Classification:
Equivalents:

Abstract

PROBLEM TO BE SOLVED: To enable to effectively obtain the subject compound useful as an intermediate for synthesizing antifogging agents for heat-developable photosensitizing materials, etc., at a low cost by selectively reducing a nitrophenylacetylene compound with iron (compound).

SOLUTION: This method for producing an aminophenylacetylene compound of formula II comprises selectively reducing (A) a compound of formula I [$R<1>$ is H, a group of the formula: $CR<2>R<3>OH$ ($R<2>$, $R<3>$ are each H, an alkyl, or $R<2>$ and $R<3>$ may be combined with each other to form a five to seven-membered ring), a group of the formula: $SiR<4>R<5>R<6>$ ($R<4>$ to $R<6>$ are each an alkyl)] with (B) iron (salt) (e.g. iron powder or reduced iron activated with acetic acid, hydrochloric acid, ammonium chloride or a nickel chloride, the mixture of ferric trichloride with a hydrogen-donor such as a hydrazine compound, ferrous dichloride or ferric trichloride). The reaction is preferably carried out by reacting 1 equivalent of the component A with 0.1-10 equivalents of the component B at a temperature of 0-150 deg.C.

Data supplied from the esp@cenet database - 12

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 11

ACID ADDUCT SALT OF 3-ETHYNYLANILINE COMPOUND AND PURIFICATION OF 3-ETHYNYLANILINE COMPOUND

Patent Number: JP10036326
Publication date: 1998-02-10
Inventor(s): YAMAKAWA KAZUYOSHI; SATO TADAHISA
Applicant(s): FUJI PHOTO FILM CO LTD
Requested Patent: ☐ JP10036326
Application Number: JP19960207787 19960718
Priority Number(s):
IPC Classification: C07C211/46; C07C209/84
EC Classification:
Equivalents:

Abstract

PROBLEM TO BE SOLVED: To obtain the subject new acid adduct salt having the form of a specific acid adduct salt, capable of being easily crystallized for its purification, excellent in storage stability and useful as an intermediate for synthesizing thermosetting resins, nonlinear optical materials, etc.

SOLUTION: A compound of formula I [$X \leftrightarrow$ is $BF_4 \leftrightarrow$, $PF_6 \leftrightarrow$, $ClO_4 \leftrightarrow$, a halogen ion, a group of formula II ($R < 1 >$ is OH, an alkyl, an aryl), a group of formula III (Z is a single bond, methylene, ethylene, phenylene)]. For example, 3-ethynylaniline sulfuric acid salt. The compound of formula I is obtained by dissolving a 3-ethynylaniline compound in an organic solvent (preferably an aromatic hydrocarbon solvent, an aliphatic hydrocarbon solvent, an ester solvent), adding an acid of the formula: HX to the solution and subsequently filtering off the deposited crystals.

Data supplied from the esp@cenet database - I2

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 12

QUINAZOLINE DERIVATIVE

Patent Number: EP0726267

Publication date: 1996-08-14

Inventor(s): MORIYAMA TAKAHIRO (JP); NONAKA HIROMI (JP); KARASAWA AKIRA (JP); OKAMURA YUKO (JP); TAKAI HARUKI (JP); YAO KOZO (JP); FUJIWARA SHIGEKI (JP)

Applicant(s): KYOWA HAKKO KOGYO KK (JP)

Requested Patent: ☐ EP0726267, A4, B1

Application Number: EP19950929231 19950825

Priority Number (s): WO1995JP01694 19950825; JP19940202018 19940826

IPC Classification: C07D401/14; A61K31/55; A61K31/505

EC Classification: C07D401/14, C07D401/14R

Equivalents: AU3265595, AU689304, CA2174854, CN1043991B, CN1134150, DE69519469D, DE69519469T, ES2153491T, FI961758, ☐ JP9165385, NO310658B, NO961601, NZ291506, ☐ US5948784, ☐ WO9606841

Cited Documents:

Abstract

Disclosed are quinazoline derivatives represented by formula (I): wherein R<1> represents hydrogen, lower alkyl, alkenyl, or aralkyl; R<2>, R<3>, R<4>, and R<5> represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyl, or the like; R<6>, R<7>, R<8>, and R<9> represent hydrogen, lower alkyl, lower alkoxy, aralkyloxy, or the like, or any adjoining two of them are combined to form methylenedioxy or the like; R<10> represents hydrogen, lower alkyl, or the like; R<11> and R<12> represent hydrogen, lower alkyl, cycloalkyl, phenyl, or aralkyl, or R<11> and R<12> are combined together with N to form a heterocyclic group; and n represents 0, 1 or 2, and pharmaceutically acceptable salts thereof. These compounds have adenosine uptake inhibitory activity and are useful for the protection of myocardium and for the prevention or treatment of inflammation such as leg and foot edema.

Data supplied from the esp@cenet database - 12

Quinazoline derivatives.Patent Number: ☐ EP0566226, B1Publication
date: 1993-10-20

Inventor(s): BARKER ANDREW JOHN (GB)

Applicant(s): ZENECA LTD (GB)

Requested
Patent: ☐ RU2127263Application
Number: EP19930300270 19930115Priority Number
(s): GB19920001095 19920120; GB19920013572 19920626; GB19920023735 19921112IPC
Classification: C07D239/94; C07D491/056; C07D403/12; A61K31/505EC
Classification: C07D239/94, C07D403/04, C07D491/04

Equivalents: AU3101093, AU661533, CA2086968, CZ9300043, DE69300754D, DE69300754T, ES2078798T, FI930208, HK36497, HU63153, HU9500185, IL104479, KR229294, NO301541B, NO930178, NZ245662, SK1693

Cited
Documents: GB2160201; US3985749; GB2033894; WO9214716; EP0520722**Abstract**


The invention concerns quinazoline derivatives of the formula I wherein m is 1, 2 or 3 and each R<1> includes hydroxy, amino, carboxy, carbamoyl, ureido, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, hydroxyamino, (1-4C)alkoxyamino, (2-4C)alkanoyloxyamino, trifluoromethoxy, (1-4C)alkyl, (1-4C)alkoxy and (1-3C)alkylenedioxy; n is 1 or 2 and each R<2> includes hydrogen, hydroxy, halogeno, trifluoromethyl, amino, nitro, cyano and (1-4C)alkyl; or a pharmaceutically-acceptable salt thereof; processes for their preparation; pharmaceutical compositions containing them; and the use of the receptor tyrosine kinase inhibitory properties of the compounds in the treatment of cancer.

Data supplied from the esp@cenet database - 12

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 14

Patent abridgement 245662

7) Described are the compounds



and pharmaceutically acceptable salts thereof, which are useful in the treatment of cancer.

in these compounds,

n is 1, 2 or 3;

m is 1 or 2; each

R¹ is independently OH, amino, substituted amino, carboxy, ureido, 3-phenylureido, carbamoyl, C₁-₄-alkoxycarbonyl, C₁-₄-alkylcarbamoyl, N,N-di-C₁-₄-alkylcarbamoyl, OCF₃ optionally substituted C₁-₄-alkoxy, C₁-₄-alkylthio, C₁-₄-alkylsulphinyl, C₁-₄-alkylsulphonyl, optionally substituted C₁-₄-alkyl, C₂-₄-alkanoyloxy, hydroxy-C₂-₆-alkanoyloxy, C₁-₄-alkoxy-C₂-₄-alkanoyloxy, substituted C₁-₄-alkylamino, optionally substituted benzamido, optionally substituted benzenesulphonamido, pyrrolidin-1-yl, piperidino, morpholino, piperazin-1-yl, 4-C₁-₄-alkylpiperazin-1-yl, 2-dioxopyrrolidin-1-yl or 2,5-dioxopyrrolidin-1-yl, or two R¹ groups together form a C₁-₃-alkylenedioxy group; and each R² is independently H, OH, CF₃, halo, amino, NO₂, CN, C₁-₄-alkyl, C₁-₄-alkoxy, mono- or di-C₁-₄-alkylamino, C₁-₄-alkylthio, C₁-₄-alkylsulphinyl, C₁-₄-alkylsulphonyl, C₂-₄-alkanoylamino, optionally substituted benzamido or C₂-₄-alkanoyl.

A Simple and Economical Synthetic Route to *p*-Ethynylaniline and Ethynyl-Terminated Substrates

Anastasi s P. Mellissaris¹ and Morton H. Litt¹


Department of Macromolecular Science, Case Western Reserve University, Cleveland, Ohio 44106

Received July 12, 1993 (Revised Manuscript Received June 20, 1994)

Introduction

Acetylenic compounds have been used for the synthesis of high performance polymers and for systems which exhibit nonlinear optical properties. Classical methods for the synthesis of terminal arylacetylenes in general involve manipulation of preformed, two-carbon side chains and include methods such as the Vilsmeier method¹⁻³ or the halogenation-dehydrohalogenation sequence of vinyl aromatics⁴ and ketones.^{5,6} An innovation in the synthesis of arylacetylenic compounds has been to use protecting groups.⁷ Acetylene, protected at one end, can be added to an aromatic nucleus via coupling at the free end. Subsequent removal of the protecting group generates a terminal arylacetylene. The widely accepted procedure for the addition of an acetylenic substituent to an aromatic nucleus is the Stephens-Castro coupling reaction⁸⁻¹⁰ between an aryl iodide and a protected acetylide in pyridine at reflux. More recent advances in the synthesis of arylacetylenes^{11,12} use a two-step route; the first step involves the coupling of an aryl iodide with (trimethylsilyl)acetylene (TMSA) in the presence of Pd(0)/Cu(I) in pyridine. The second step is removal of the protecting group (trimethylsilyl) to yield the arylacetylene. The trimethylsilyl protecting group is easily removed by treatment with dilute potassium hydroxide or potassium carbonate. However, because of the prohibitively high cost of the TMSA, this route has been limited to small-scale preparations. There has been a great interest in the development of methods for introducing an ethynyl group¹³⁻¹⁵ into organic structures.

Scheme 1. Synthetic Route to *p*-Ethynylaniline



For the synthesis of *p*-ethynylaniline (1c) (Scheme 1), four methods^{13,16-18} have been reported. The yields vary from poor to moderate (30–65%) and the reactions are cumbersome and costly to perform on a large scale. The most interesting procedure for the synthesis of 1c¹¹ entails coupling of *p*-iodoaniline with (trimethylsilyl)acetylene (TMSA) in the presence of a palladium complex and a copper(I) salt. Due to the high cost of TMSA, this route for all practical purposes has been limited to small-scale procedures.¹⁹⁻²³ J. Stille and T. Takeichi¹⁷ synthesized 1c using (tributylstannyl)acetylene (TBSA) and *p*-iodoaniline in 30% overall yield.

Attempts to synthesize larger quantities of 1c using inexpensive reagents have been unsuccessful up to date. 2-Methyl-3-butyn-2-ol (MEBYNOL) has been used by other investigators to synthesize 1c because of its very low cost. Bardamova et al.¹⁸ synthesized 1c on a milligram scale by direct coupling of *p*-iodoaniline with MEBYNOL, followed by deprotection and heating the intermediate 4-anilino-2-methyl-3-butyn-2-ol under a high vacuum in the presence of well-ground KOH and catalytic amounts of hydroquinone. However, most of the desired product decomposed under these severe conditions. Takalo et al.¹⁶ reported a modified procedure for deprotecting 4-anilino-2-methyl-3-butyn-2-ol, heating under distillation conditions in the presence of NaOH pellets in toluene for 2 h. 1c was synthesized in 30% overall yield. The methods of Bardamova¹⁸ and Takalo¹⁶ have not been used for the synthesis of 1c because the yields were low and some decomposition products were generated during the deprotection step.

Due to the high cost of TMSA, we decided to develop a simple high yield route to 1c using the very inexpensive reagent MEBYNOL. We have reported a new synthesis of *p*-ethynylbenzoic acid and *p*-ethynyl benzoyl chloride, using MEBYNOL.¹⁴ We now report an economical and efficient synthesis of 1c using a modified route which is simpler and less expensive than the methods previously reported. This method gives an almost quantitative yield of high purity product. The low yields and the various

¹ Current address: Materials Engineering Division, Research Institute, University of Dayton, Dayton, OH 45469.

(1) Staab, H.; Neunhoffer, K. *Synthesis* 1974, 424.
(2) Rosenblum, M.; Braun, N.; Papenmeier, J.; Applebaum, M. J. *Organomet. Chem.* 1968, 8, 173.

(3) Boschan, R. Hughes Aircraft Co. Private communication cited in ref 12.

(4) Collier, W.; Macomber, R. J. *Org. Chem.* 1973, 38, 1367.

(5) Kocienski, P. J. *Org. Chem.* 1974, 39, 3285.

(6) Bartlett, P.; Rosen, L. J. *Am. Chem. Soc.* 1942, 64, 543.

(7) Walton, D. In *Protective Groups in Organic Chemistry*; McOmie, J., Ed.; Plenum Press: London, 1973; Chapter 1.

(8) Castro, C.; Stephens, R. J. *Org. Chem.* 1963, 28, 2163.

(9) Stephens, R.; Castro, C. J. *Org. Chem.* 1963, 28, 3313.

(10) Castro, C.; Havlin, R.; Honwad, V.; Malte, A.; Moja, S. J. *Am. Chem. Soc.* 1969, 91, 6464.

(11) Takahashi, S.; Kuroyama, Y.; Sonogashira, K.; Hagihara, N. *Synthesis*, 1980, 627.

(12) Austin, W.; Billow, N.; Kallaghan, W.; Lau, K. J. *Org. Chem.* 1981, 46, 2280.

(13) Sabourin, E.; Onopchenko, A. J. *Org. Chem.* 1983, 48, 5185.

(14) Mellissaris, A.; Litt, M. J. *Org. Chem.* 1992, 57, 6998.

(15) Bardamova, M.; Trotsenko, Z.; Kotlyarskii, I. *Bull. Acad. Science USSR, Div. Chem.* 1983, 1057.

(16) Takalo, H.; Kankari, J.; Hanninen, E. *Acta Chem. Scand.* 1988, B(42), 448.

(17) Takeichi, T.; Stille, J. *Macromolecules* 1986, 19, 2093.

(18) Fujiwara, H.; Kihara, K. Japanese Patent 03 38 551, 1991; *Chem. Abstr.* 1991, 115, 49061c.

(19) Abraham, T.; Soloski, E.; Benner, C. L.; Evers, R. C. Report 1989, WRDC-TR-89-4116, 1990, 90(12); *Chem. Abstr.* 1991, 116, 8379c.

(20) Yuan, Z.; Taylor, N. J.; Marder, T. B.; Williams, I. D.; Kurtz, S. K.; Cheng, L. T. J. *Chem. Soc., Chem. Commun.* 1990, 1489.

(21) Schering, A. G. Ger. Offen. DE 3 818 062, 1989; *Chem. Abstr.* 1990, 113, 40867f.

(22) Choe, E. W. U.S. Patent 4,703,096, 1987; *Chem. Abstr.* 1988, 108, 76103a.

(23) Graham, E. M.; Miskowski, V. M.; et al. *J. Am. Chem. Soc.* 1989, 111(24), 8771.

XP-000916169

D2

FG18

J. Org. Chem. 1994, 59, 5818-5821

A Simple and Economical Synthetic Route to *p*-Ethynylaniline and Ethynyl-Terminated Substrates

Anastasi s P. Melissaris[†] and Morton H. Litt^{*}


Department of Macromolecular Science, Case Western Reserve University, Cleveland, Ohio 44106

Received July 12, 1993 (Revised Manuscript Received June 20, 1994)

Introduction

Acetylenic compounds have been used for the synthesis of high performance polymers and for systems which exhibit nonlinear optical properties. Classical methods for the synthesis of terminal arylacetylenes in general involve manipulation of preformed, two-carbon side chains and include methods such as the Vilsmeier method¹⁻³ or the halogenation-dehydrohalogenation sequence of vinyl aromatics⁴ and ketones.^{5,6} An innovation in the synthesis of arylacetylenic compounds has been to use protecting groups.⁷ Acetylene, protected at one end, can be added to an aromatic nucleus via coupling at the free end. Subsequent removal of the protecting group generates a terminal arylacetylene. The widely accepted procedure for the addition of an acetylenic substituent to an aromatic nucleus is the Stephens-Castro coupling reaction⁸⁻¹⁰ between an aryl iodide and a protected acetylide in pyridine at reflux. More recent advances in the synthesis of arylacetylenes^{11,12} use a two-step route; the first step involves the coupling of an aryl iodide with (trimethylsilyl)acetylene (TMSA) in the presence of Pd(0)/Cu(I) in pyridine. The second step is removal of the protecting group (trimethylsilyl) to yield the arylacetylene. The trimethylsilyl protecting group is easily removed by treatment with dilute potassium hydroxide or potassium carbonate. However, because of the prohibitively high cost of the TMSA, this route has been limited to small-scale preparations. There has been a great interest in the development of methods for introducing an ethynyl group¹³⁻¹⁸ into organic structures.

Scheme 1. Synthetic Route to *p*-Ethynylaniline



For the synthesis of *p*-ethynylaniline (1c) (Scheme 1), four methods^{13,16-18} have been reported. The yields vary from poor to moderate (30–65%) and the reactions are cumbersome and costly to perform on a large scale. The most interesting procedure for the synthesis of 1c¹¹ entails coupling of *p*-iodoaniline with (trimethylsilyl)acetylene (TMSA) in the presence of a palladium complex and a copper(I) salt. Due to the high cost of TMSA, this route for all practical purposes has been limited to small-scale procedures.¹⁹⁻²² J. Stille and T. Takeichi¹⁷ synthesized 1c using (tributylstannyl)acetylene (TBSA) and *p*-iodoaniline in 30% overall yield.

Attempts to synthesize larger quantities of 1c using inexpensive reagents have been unsuccessful up to date. 2-Methyl-3-butyne-2-ol (MEBYNOL) has been used by other investigators to synthesize 1c because of its very low cost. Bardamova et al.²³ synthesized 1c on a milligram scale by direct coupling of *p*-iodoaniline with MEBYNOL, followed by deprotection and heating the intermediate 4-anilino-2-methyl-3-butyne-2-ol under a high vacuum in the presence of well-ground KOH and catalytic amounts of hydroquinone. However, most of the desired product decomposed under these severe conditions. Takalo et al.¹⁶ reported a modified procedure for deprotecting 4-anilino-2-methyl-3-butyne-2-ol, heating under distillation conditions in the presence of NaOH pellets in toluene for 2 h. 1c was synthesized in 30% overall yield. The methods of Bardamova²³ and Takalo¹⁶ have not been used for the synthesis of 1c because the yields were low and some decomposition products were generated during the deprotection step.

Due to the high cost of TMSA, we decided to develop a simple high yield route to 1c using the very inexpensive reagent MEBYNOL. We have reported a new synthesis of *p*-ethynylbenzoic acid and *p*-ethynyl benzoyl chloride, using MEBYNOL.¹⁴ We now report an economical and efficient synthesis of 1c using a modified route which is simpler and less expensive than the methods previously reported. This method gives an almost quantitative yield of high purity product. The low yields and the various

[†] Current address: Materials Engineering Division, Research Institute, University of Dayton, Dayton, OH 45469.

- (1) Staab, H.; Neunhoffer, K. *Synthesis* 1974, 424.
- (2) Rosenblum, M.; Brown, N.; Papenmeier, J.; Applebaum, M. J. *Organomet. Chem.* 1968, 8, 173.
- (3) Boschan, R. Hughes Aircraft Co. Private communication cited in ref 12.
- (4) Collier, W.; Macomber, R. J. *Org. Chem.* 1973, 38, 1367.
- (5) Kocienski, P. J. *Org. Chem.* 1974, 39, 3285.
- (6) Bartlett, P.; Rosen, L. J. *Am. Chem. Soc.* 1942, 64, 543.
- (7) Walton, D. In *Protective Groups in Organic Chemistry*; McOmie, J., Ed.; Plenum Press: London, 1973; Chapter 1.
- (8) Castro, C.; Stephens, R. J. *Org. Chem.* 1963, 28, 2163.
- (9) Stephens, R.; Castro, C. J. *Org. Chem.* 1963, 28, 3313.
- (10) Castro, C.; Havlin, R.; Howard, V.; Malta, A.; Moja, S. J. *Am. Chem. Soc.* 1969, 91, 6464.
- (11) Takahashi, S.; Kuroyama, Y.; Sonogashira, K.; Hagihara, N. *Synthesis*, 1980, 627.
- (12) Austin, W.; Billow, N.; Kallaghan, W.; Lau, K. J. *Org. Chem.* 1981, 46, 2280.
- (13) Sabourin, E.; Onopchenko, A. J. *Org. Chem.* 1988, 49, 5135.
- (14) Melissaris, A.; Litt, M. J. *Org. Chem.* 1992, 57, 6998.
- (15) Bardamova, M.; Trotsenko, Z.; Kotlyarskii, I. *Bull. Akad. Sci. USSR, Div. Chem.* 1983, 1067.
- (16) Takalo, H.; Kankari, J.; Hanninen, E. *Acta Chem. Scand.* 1988, B(42), 448.
- (17) Takeichi, T.; Stille, J. *Macromolecules* 1986, 19, 2093.
- (18) Fujiwara, H.; Kihara, K. Japanese Patent 03 38 551, 1991; *Chem. Abstr.* 1991, 115, 49061c.

- (19) Abraham, T.; Soloski, E.; Benner, C. L.; Evans, R. C. *Report 1989, WRDC-TR-89-4115*, 1990, 90(12); *Chem. Abstr.* 1991, 115, 9379c.
- (20) Yuan, Z.; Taylor, N. J.; Marder, T. B.; Williams, I. D.; Kurtz, S. K.; Cheng, L. T. J. *Chem. Soc., Chem. Commun.* 1990, 1489.
- (21) Schering, A. G. Ger. Offen. DE 3 818 062, 1989; *Chem. Abstr.* 1990, 113, 40867f.
- (22) Choe, E. W. U.S. Patent 4,703,096, 1987; *Chem. Abstr.* 1988, 108, 76103a.
- (23) Graham, E. M.; Miskowski, V. M.; et al. *J. Am. Chem. Soc.* 1989, 111(24), 8771.

LAWYERS' AND MERCHANTS' TRANSLATION BUREAU, INC.

Legal, Financial, Scientific, Technical and Patent Translations

11 BROADWAY

NEW YORK, NY 10004



Certificate of Accuracy

TRANSLATION

From Spanish into English

**STATE OF NEW YORK } s.s.:
COUNTY OF NEW YORK }**

On this day personally appeared before me
who, after being duly sworn, deposes and states: Elisabeth A. Lucas

That he is a translator of the Spanish and English languages by profession and
as such connected with the **LAWYERS' & MERCHANTS' TRANSLATION
BUREAU;**

That he is thoroughly conversant with these languages;

That he has carefully made the attached translation from the original document
written in the Spanish language; and

That the attached translation is a true and correct English version of such original,
to the best of his knowledge and belief.

**SUBSCRIBED AND SWORN TO BEFORE ME
THIS**


SEP 16 2003

**Susan Tapley
Notary Public, State of New York
No. 01TA4999804
Qualified in Queens County
Certificate filed in New York County
and Kings County
Commission Expires July 27, 2006**

**Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 18**


was prepd. in 5 steps from 6-methyl-5-nitroindazole. Key intermediate was the aminoindazolecarboxylic acid III, which was annulated by heating at 70° with $\text{HC}(\text{NH})\text{NH}_2\text{AcOH}$ for 2 days to give 91% IL I ($\text{R}^1 = \text{O}$, $\text{R}^2 = \text{H}$) was prepd. in 76% yield from III by fusion with $(\text{H}_2\text{N})_2\text{CO}$ at 160° for 15 min. Pyrazoloquinazolinone IV was prepd. in 35% overall yield from 5,2-Me(HO_2C) $\text{C}_6\text{H}_4\text{NHAc}$ in 8 steps and pyrazoloquinazolinone V was prepd., in 5 steps, from 6-methyl-4-nitroindazole.

98: 122723r Synthesis of 2- and 4-iminopyrido[1,2-a]pyrimidines from allenic nitriles and 2-aminopyridines. Fomum, Z. Tanee; Mbafor, J. Tanyi; Landor, Phyllis D.; Landor, Stephen R. (Univ. Yaounde, Yaounde, Cameroon). *Tetrahedron Lett.* 1981, 22(41), 4127-8 (Eng). Heating $\text{RCR}^1\text{C}:\text{C}:\text{CHCN}$ (I; R




= Me, $\text{R}^1 = \text{Et}$, Pr; R = $\text{R}^1 = \text{Et}$) under reflux with 2-amino- or 2,3-diaminopyridine in alc. soln. for 48 h gave 87-94% 4-imino- or 2-imino-2,3-diaminopyridine II (R, R^1 as before, $\text{R}^2 = \text{H}$, NH_2 , resp.), whereas similar treatment of I ($\text{R} = \text{R}^1 = \text{Me}$; R = Pr, $\text{R}^1 = \text{H}$) with 2-aminopyridine and I ($\text{R} = \text{Me}$, Et, $\text{R}^1 = \text{Et}$) with 3-hydroxy-2-aminopyridine gave 90-2% 2-iminopyrido[1,2-a]pyrimidine III (R, R^1 as before, $\text{R}^2 = \text{H}$, OH, resp.).

98: 122724s Synthesis of 6-alkoxy-2-amino-5-cyanopyrimidines through sodium alkoxide-induced regioselective cyclization of 1,3-dicarbonyltrifiles. Perez, Miguel A.; Soto, Jose L. (Fac. Quim., Univ. Complutense, Madrid, Spain). *Synthesis* 1981, (12), 955-8 (Eng). The reaction of $\text{R}^1\text{C}(\text{OR})\text{C}(\text{CN})\text{R}^2$




(R = Et, Me; $\text{R}^1 = \text{H}$, Me, Ph, tolyl, anisyl, ClC_6H_4 , $\text{O}_2\text{NC}_6\text{H}_4$) with $\text{H}_2\text{N-CN}$ and $\text{NaOR}^2\text{-R}^2\text{OH}$ ($\text{R}^2 = \text{Me}$, Et, Pr) yielded the resp. pyrimidinetrifiles I. Similarly, pyrimidines II were prepd. from $\text{R}^1\text{C}(\text{OR})\text{C}(\text{CN})\text{R}^2$, $\text{H}_2\text{N}(\text{OMe})\text{N}:\text{H}:\text{Cl}$, and NaOMe-MeOH . A mixt. of $\text{EtOCH}_2\text{C}(\text{CN})_2$, H_2NCH_3 , and NaOMe in MeOH was refluxed to give I ($\text{R}^1 = \text{H}$, $\text{R}^2 = \text{Me}$).

98: 122725t Preparation of 3-aryl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazolinones from methyl N-aryldithiocarbamates and anthranilic acid. Mayoral, J.; Melendez, E.; Merchan, F.; Sanchez, J. (Dep. Quim. Org., Univ. Zaragoza, Zaragoza, Spain). *Synthesis* 1981, (12), 962 (Eng). The cyclocondensation




reaction of RNHCS_2Me (R = tolyl, xylyl, methylchlorophenyl, (methylenedioxy)phenyl (trifluoromethyl)phenyl, Ph, anisyl) with anthranilic acids gave the resp. quinazolinethiones I ($\text{R}^1 = \text{H}$, Cl; $\text{R}^2 = \text{H}$, NO₂, Cl; $\text{R}^3 = \text{H}$, Cl). Thus, 4-MeC₆H₄NHCS₂Me in DMF was added dropwise to 2-H₂N-C₆H₄-CO₂H in DMF at room temp., and the mixt. was refluxed to give I (R = 4-MeC₆H₄, $\text{R}^1 = \text{R}^2 = \text{R}^3 = \text{H}$).

98: 122726u Studies on antimalarial agents. VI. Synthesis and their antimalarial activities of 2,4-diamino-6-substituted-aminosulfonylquinazolinone derivatives. Zhang, Xiping; Shen, Defu; Zhang, Xijun; Chen, Lin; Dai, Zuru; Shu, Kangqun (Shanghai Inst. Pharm. Ind. Res., Shanghai, Peop. Rep. China). *Yaoxue Xuebao* 1981, 16(11), 877-80 (Ch).




Sixteen quinazolinonesulfonamides I [$\text{NRR}^1 = \text{alkylamino}$, 1-pyrrolidinyl, piperidinyl (II), morpholino, etc.] were prepd. by amidation of 2,4-diaminoquinazolinone-6-sulfonyl chloride. II showed pronounced prophylactic activity against *Plasmodium yoelii*.

98: 122727v Studies on drugs for coronary diseases. II. Synthesis of compounds related to changrolin, a new antiarrhythmic agent. Sun, Cunji; Zhang, Xinyi; Yang, Kingzhong; Wang, Pingping; Shen, Jian; Shu, Yun; Ji, Ruyun; Kyl, Zuyong (Shanghai Inst. Mater. Med., Acad. Sin., Shanghai, Peop. Rep. China). *Yaoxue Xuebao* 1981, 16(8), 564-70 (Ch). Changrolin (I; $\text{NR}_2 = 1\text{-pyrrolidinyl}$, $\text{R}^1 = 1\text{-pyrrolidinylmethyl}$, $\text{R}^2 = \text{H}$, X = N) analogs, i.e., I [$\text{NR}_2 = \text{NMe}_2$, $\text{N}(\text{CH}_3)_2\text{CH}_2\text{OH}$], morpholino, etc.; $\text{R}^1 = \text{H}$, CH_2NR_2 ; $\text{R}^2 = \text{H}$; X = N], I [$\text{NR}_2 = \text{NMe}_2$, 1-pyrrolidinyl; $\text{R}^1 = \text{H}$, CH_2NR_2 ; $\text{R}^2 = \text{Cl}$; X = CH], II




($\text{NZ} = \text{NMe}_2$, 1-pyrrolidinyl, 1-piperidinyl, morpholino), III ($\text{NZ} = \text{NMe}_2$, 1-pyrrolidinyl, morpholino, etc.; $\text{R}^2 = \text{Ac}$, Bz; $\text{R}^4 = \text{H}$, CH_2NZ) and IV ($\text{NZ} = \text{NMe}_2$, 1-pyrrolidinyl, morpholino, etc.) were prepd. by known reactions. III ($\text{NZ} = 1\text{-pyrrolidinyl}$, $\text{R}^2 = \text{Bz}$, $\text{R}^4 = \text{CH}_2\text{NZ}$) was more effective than changrolin in protecting dogs from arterial fibrillation.

98: 122728w Studies on antiarrhythmics - synthesis of 2-[(alkylamino)methyl]- and 2,6-bis[(alkylamino)methyl]-4-[(substituted amino)phenols]. Lin, Mulan; Liu, Yufang; Lu, Yongyu; Zhang, Huiqin; Zheng, Weimin (Tianjin Inst. Pharm. Ind. Res., Tianjin, Peop. Rep. China). *Yaoxue Xuebao* 1981, 16(10), 757-61 (Ch). 4-Aminophenols I [R = Ac, 2,6-diamino(or




dimethyl)-4-pyrimidinyl, 1-phthalazinyl, 6,7-dimethoxy-4-quinazolinyl, etc.; $\text{R}^1, \text{R}^2 = \text{H}$, Et , NCH_3 , 1-pyrrolidinylmethyl, piperidinomethyl, morpholinomethyl] (37 compds.) were prepd. by known reactions. Some I showed antiarrhythmic activity.

98: 122729x Redox reactions of 10H-pyrimido[5,4-b][1,4]benzothiazines. Fennel, Helmut; Roessler, Hellmuth H.; Grauert, Rolf W. (Inst. Pharm., Freien Univ., D-1000 Berlin, 33 Fed. Rep. Ger.). *Arch. Pharm. (Weinheim, Ger.)* 1981, 314(12), 1023-30 (Ger). The structure, spectra, and reactivity of mcl.



species participating in the thiazaloxazine redox system are described. Oxidn. of I ($\text{R} = \text{H}$, Me, n = 0) gave the cations II which were hydrolyzed to I (n = 1) or alcoholized to III ($\text{R}^1 = \text{H}$, Me, Et, Pr, CHMe, Bu).

98: 122730r Reaction of 1,2,3,4-tetrahydroquinazolin-4-ones with acid anhydride. III. Yamato, Masatoshi; Horuchi, Jiro; Takeuchi, Yasuo (Fac. Pharm. Sci., Okayama Univ., Okayama, Japan 700). *Chem. Pharm. Bull.* 1981, 29(11), 3124-9 (Eng). The reaction of C₆-substituted 1,2,3,4-tetrahy-



droquinazolin-4-ones with Ac_2O and pyridine was carried out in order to elucidate the effect of the C₆-substituent. It was found that the various types of reactions occurred depending on the kind and no. of C₆-substituents of 1,2,3,4-tetrahydroquinazolin-4-ones. Thus, the quinazolinone I (R = PhCH₂CH₃) was treated with Ac_2O at 100° for 3 h to give the quinazolinone II (21%). I (R = Ph) reacted with Ac_2O to give 68% o-(PhCMe₂N)C₆H₄CONHAc.

98: 122731s Studies on fluorinated pyrimidines. I. A new method of synthesizing 5-fluorouracil and its derivatives. Miyashita, Osamu; Matsumura, Koichi; Shimadzu, Hiroshi; Hashimoto, Naoto (Cent. Res. Div., Takeda Chem. Ind., Ltd., Osaka, Japan 532). *Chem. Pharm. Bull.* 1981, 29(11), 3181-90 (Eng). Title compds. I (R = H, $\text{R}^1 = \text{F}$, $\text{R}^2 = \text{CO}_2\text{Me}$, $\text{R}^3 = \text{H}$,

Applicants: Timothy Norris et al.

Serial No.: 09/711,272

Filed: November 9, 2000

Exhibit 19



XP 000517991

Pergamon

134a Tetrahedron Letters

36(1995) August 14, No. 33, Kidlington, Oxford, GB

Tetrahedron Letters, Vol. 36, No. 33, pp. 5891-5894, 1995

Elsevier Science Ltd

Printed in Great Britain

0040-4039/95 \$9.50+0.00

p. 5891-5894 = (4)

0040-4039(95)01172-2

C67D517/72C


A Convergent Synthesis of Functionalized B-seco Taxane Skeletons

Christian Montalbetti, Monique Savignac, Félicie Bonnefis and Jean Pierre Genêt.


Laboratoire de Synthèse Organique, associé au CNRS, Ecole Nationale Supérieure de Chimie de Paris,
11 rue Pierre et Marie Curie - 75231 Paris Cedex 05 - France

Abstract: The sequential Sonogashira cross-coupling reactions with water soluble and anhydrous Pd(0) catalysts between vinylic iodo derivatives 6, 8 and 3-iodocyclohexenone 7 with trimethyl silyl acetylene are used to produce functionalized intermediates 11 and 18. Conjugate addition followed by enolate trapping with trimethyl orthoformate provided B-seco taxane derivatives 14 and 20.

The antitumor agents, paclitaxel (Taxol®) 1 and docetaxel (Taxotere®) 2 have generated much excitement due to their activities against advanced ovarian and breast cancer.¹ Taxol 1 has been the subject of extensive chemical and biological studies, which have been summarized in recent reviews.^{1c,2} The recent total syntheses of taxol accomplished by Nicolaou³ and Holton⁴ are seminal achievements in the field.



The challenge now is to provide new methodologies for the synthesis of 10-deacetylpaclitaxin III 3 2b analogues⁵ which can rapidly lead to the analogues of taxol and taxotere.



Scheme 1

We wish to report a convergent synthesis of B-seco taxane precursors of taxoids 4 by linking the future A and C rings through a two carbon moiety, via a sequential Sonogashira⁶ reaction between the protected iodo-ketone 6 and 3-iodocyclohexenone 7 (Scheme 1).

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 17

LAWYERS' AND MERCHANTS' TRANSLATION BUREAU, INC.
Legal, Financial, Scientific, Technical and Patent Translations
11 BROADWAY
NEW YORK, NY 10004



Certificate of Accuracy

TRANSLATION
From Japanese into English

STATE OF NEW YORK
COUNTY OF NEW YORK

} s.s. :

On this day personally appeared before me Elisabeth A. Lucas
who, after being duly sworn, deposes and states:

That he is a translator of the Japanese and English languages by
profession and as such connected with the **LAWYERS' & MERCHANTS'**
TRANSLATION BUREAU;

That he is thoroughly conversant with these languages;

That he has carefully made the attached translation from the original document
written in the Japanese language; and

That the attached translation is a true and correct English version of such original,
to the best of his knowledge and belief.

SUBSCRIBED AND SWORN TO BEFORE ME
THIS

SEP 11 2000
Susan Tapley
Susan Tapley
Notary Public, State of New York
No. 01TA4999804
Qualified in Queens County
Certificate filed in New York County
and Kings County
Commission Expires July 27, 2006

[Handwritten signature]

Applicants: Timothy Norris et al.
Serial No.: 09/711,272
Filed: November 9, 2000
Exhibit 10

QUINAZOLINE DERIVATIVE

Patent Number: EP0726267
 Publication date: 1996-08-14
 Inventor(s): MORIYAMA TAKAHIRO (JP); NONAKA HIROMI (JP); KARASAWA AKIRA (JP); OKAMURA YUKO (JP); TAKAI HARUKI (JP); YAO KOZO (JP); FUJIWARA SHIGEKI (JP)
 Applicant(s): KYOWA HAKKO KOGYO KK (JP)
 Requested Patent: ☐ EP0726267, A4, B1
 Application Number: EP19950929231 19950825
 Priority Number (s): WO1995JP01694 19950825; JP19940202018 19940826
 IPC Classification: C07D401/14; A61K31/55; A61K31/505
 EC Classification: C07D401/14, C07D401/14R
 Equivalents: AU3265595, AU689304, CA2174854, CN1043991B, CN1134150, DE69519469D, DE69519469T, ES2153491T, FI961758, ☐ JP9165385, NO310658B, NO961601, NZ291506, ☐ US5948784, ☐ WO9606841

Cited Documents:

Abstract

Disclosed are quinazoline derivatives represented by formula (I): wherein R<1> represents hydrogen, lower alkyl, alkenyl, or aralkyl; R<2>, R<3>, R<4>, and R<5> represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyl, or the like; R<6>, R<7>, R<8>, and R<9> represent hydrogen, lower alkyl, lower alkoxy, aralkyloxy, or the like, or any adjoining two of them are combined to form methylenedioxy or the like; R<10> represents hydrogen, lower alkyl, or the like; R<11> and R<12> represent hydrogen, lower alkyl, cycloalkyl, phenyl, or aralkyl, or R<11> and R<12> are combined together with N to form a heterocyclic group; and n represents 0, 1 or 2, and pharmaceutically acceptable salts thereof. These compounds have adenosine uptake inhibitory activity and are useful for the protection of myocardium and for the prevention or treatment of inflammation such as leg and foot edema.

Data supplied from the esp@cenet database - 12

EP0566226 [Biblio](#) [Desc](#) [Claims](#) [Page 1](#)

esp@cenet

Quinazoline derivatives.Patent Number: ☐ [EP0566226](#), [B1](#)

Publication date: 1993-10-20

Inventor(s): BARKER ANDREW JOHN (GB)

Applicant(s): ZENECA LTD (GB)

Requested Patent: ☐ [RU2127263](#)

Application Number: EP19930300270 19930115

Priority Number (s): GB19920001095 19920120; GB19920013572 19920626; GB19920023735 19921112

IPC Classification: C07D239/94; C07D491/056; C07D403/12; A61K31/505

EC Classification: [C07D239/94](#), [C07D403/04](#), [C07D491/04](#)

Equivalents: AU3101093, AU661533, CA2086968, CZ9300043, DE69300754D, DE69300754T, ES2078798T, FI930208, HK36497, HU63153, HU9500185, IL104479, KR229294, NO301541B, NO930178, NZ245662, SK1693

Cited Documents: [GB2160201](#); [US3985749](#); [GB2033894](#); [WO9214716](#); [EP0520722](#)**Abstract**


The invention concerns quinazoline derivatives of the formula I wherein m is 1, 2 or 3 and each R<1> includes hydroxy, amino, carboxy, carbamoyl, ureido, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-[(1-4C)alkyl]carbamoyl, hydroxyamino, (1-4C)alkoxyamino, (2-4C)alkanoyloxyamino, trifluoromethoxy, (1-4C)alkyl, (1-4C)alkoxy and (1-3C)alkylenedioxy; n is 1 or 2 and each R<2> includes hydrogen, hydroxy, halogeno, trifluoromethyl, amino, nitro, cyano and (1-4C)alkyl; or a pharmaceutically-acceptable salt thereof; processes for their preparation; pharmaceutical compositions containing them; and the use of the receptor tyrosine kinase inhibitory properties of the compounds in the treatment of cancer.

Data supplied from the esp@cenet database - I2

Applicants: Timothy Norris et al.
 Serial No.: 09/711,272
 Filed: November 9, 2000
 Exhibit 14

Patent abridgement 245662

7) Described are the compounds



and pharmaceutically acceptable salts thereof, which are useful in the treatment of cancer.

and these compounds,

n is 1, 2 or 3;

m is 1 or 2; each

R¹ is independently OH, amino, substituted amino, carboxy, ureido, 3-phenylureido, carbamoyl, C₁-₄-alkoxycarbonyl, C₁-₄-alkylcarbamoyl, N,N-di-C₁-₄-alkylcarbamoyl, OCF₃ optionally substituted C₁-₄-alkoxy, C₁-₄-alkylthio, C₁-₄-alkylsulphanyl, C₁-₄-alkylsulphonyl, optionally substituted C₁-₄-alkyl, C₂-₄-alkanoyloxy, hydroxy-C₂-₆-alkanoyloxy, C₁-₄-alkoxy-C₂-₄-alkanoyloxy, substituted C₁-₄-alkylamino, optionally substituted benzamido, optionally substituted benzenesulphonamido, pyrrolidin-1-yl, piperidino, morpholino, piperazin-1-yl, 4-C₁-₄-alkylpiperazin-1-yl, 2-pyrrolidin-1-yl or 2,5-dioxopyrrolidin-1-yl, or two R¹ groups together form a C₁-₃-alkylenedioxy group; and each R² is independently H, OH, CF₃, halo, amino, NO₂, CN, C₁-₄-alkyl, C₁-₄-alkoxy, mono- or di-C₁-₄-alkylamino, C₁-₄-alkylthio, C₁-₄-alkylsulphanyl, C₁-₄-alkylsulphonyl, C₂-₄-alkanoylamino, optionally substituted benzamido or C₂-₄-alkanoyl.